

# Almost Engel compact groups

Evgeny Khukhro

University of Lincoln, UK

Ischia, March 2018

Joint work with Pavel Shumyatsky

# Engel groups

Notation: left-normed simple commutators

$$[a_1, a_2, a_3, \dots, a_r] = [\dots[[a_1, a_2], a_3], \dots, a_r].$$

Recall: a group  $G$  is an **Engel group** if for every  $x, g \in G$ ,

$$[x, g, g, \dots, g] = 1,$$

where  $g$  is repeated sufficiently many times depending on  $x$  and  $g$ .

Clearly, any locally nilpotent group is an Engel group.

# Known facts on finite groups

## Zorn's Theorem

*A finite Engel group is nilpotent.*

### **Proof:**

Coprime action  $\Rightarrow$  non-Engel.

No coprime action  $\Rightarrow$  nilpotent. □

## Baer's Theorem

*If  $g$  is an Engel element of a finite group  $G$ , that is,  $[x, g, \dots, g] = 1$  for every  $x \in G$ , then  $g \in F(G)$ .*

Here,  $F(G)$  is the Fitting subgroup, largest normal nilpotent subgroup.

## Engel compact groups

J. Wilson and E. Zelmanov, 1992

*Any Engel profinite group is locally nilpotent.*

Proof relies on

### Zel'manov's Theorem

*If a Lie algebra  $L$  satisfies a nontrivial identity and is generated by  $d$  elements such that each commutator in these generators is ad-nilpotent, then  $L$  is nilpotent.*

Yu. Medvedev, 2003

*Any Engel compact (Hausdorff) group is locally nilpotent.*

# Almost Engel groups

## Definition

A group  $G$  is **almost Engel** if for every  $g \in G$  there is a finite set  $\mathcal{E}(g)$  such that for every  $x \in G$ ,

$$[x, \underbrace{g, g, \dots, g}_n] \in \mathcal{E}(g) \quad \text{for all } n \geq n(x, g).$$

Includes Engel groups: when  $\mathcal{E}(g) = \{1\}$  for all  $g \in G$ .

## Theorem 1 (Almost Engel $\Rightarrow$ almost locally nilpotent)

*Suppose that  $G$  is an almost Engel compact (Hausdorff) group. Then  $G$  has a finite normal subgroup  $N$  such that  $G/N$  is locally nilpotent.*

(...there is also a locally nilpotent subgroup of finite index:  $C_G(N)$ .)

# Three parts of the proof

1. **Finite groups**, a quantitative version.
2. **Profinite groups**: using finite groups, Wilson–Zelmanov theorem.
3. **Compact groups**: reduction to profinite case using structure theorems for compact groups.

## Some notation

If  $G$  is an almost Engel group, then for every  $g \in G$  there is a unique **minimal** finite set  $\mathcal{E}(g)$  with the property that for every  $x \in G$ ,

$$[x, \underbrace{g, g, \dots, g}_n] \in \mathcal{E}(g) \quad \text{for all } n \geq n(x, g)$$

(for possibly larger numbers  $n(x, g)$ ).

We fix the symbols  $\mathcal{E}(g)$  for these minimal sets, call them **Engel sinks**.

The **nilpotent residual** of a group  $G$  is

$$\gamma_\infty(G) = \bigcap_i \gamma_i(G),$$

where  $\gamma_i(G)$  are terms of the lower central series ( $\gamma_1(G) = G$ , and  $\gamma_{i+1}(G) = [\gamma_i(G), G]$ ).

# Almost Engel finite groups

For finite groups there must be a quantitative analogue of the hypothesis that the sinks  $\mathcal{E}(g)$  are finite.

## Theorem 2

*Suppose that  $G$  is a finite group and there is a positive integer  $m$  such that  $|\mathcal{E}(g)| \leq m$  for every  $g \in G$ . Then  $|\gamma_\infty(G)|$  is bounded in terms of  $m$ .*

(...And  $G$  also has a nilpotent normal subgroup of bounded index:  
 $C_G(\gamma_\infty(G))$ .)

## About the proof for finite groups

### Lemma

In any almost Engel group  $G$ , the Engel sink is the set

$$\mathcal{E}(g) = \{z \in G \mid z = [z, g, \dots, g]\}$$

(with at least one occurrence of  $g$ ).

Indeed,  $x \rightarrow [x, g]$  is a mapping of  $\mathcal{E}(g)$  into itself,

must be “onto” since  $\mathcal{E}(g)$  is finite and minimal,

$z \in \mathcal{E}(g)$  belongs to its orbit. □

### Lemma

*In a finite group, if  $A$  is an abelian section, acted on by  $g$  of coprime order, then  $[A, g] = \{[a, g, \dots, g] \mid a \in A\}$  for any number of  $g$ , so  $[A, g] \subseteq \mathcal{E}(g)$ .*

**Proof:**  $C_{[A, g]}(g) = 1 \Rightarrow [A, g] = \{[b, g] \mid b \in [A, g]\}$ . □

## About the proof for finite groups

### Lemma

If  $|\mathcal{E}(g)| \leq m$  for all  $g \in G$ , then  $G/F(G)$  is of  $m$ -bounded exponent.

**Proof:** Clearly,  $g$  centralizes its powers. Hence for any  $z \in \mathcal{E}(g^k)$  we have

$$z = [z, g^k, \dots, g^k] \Rightarrow z^g = [z^g, g^k, \dots, g^k].$$

Therefore  $\mathcal{E}(g^k)$  is  $g$ -invariant.

Choose  $k = m!$ . Then  $g^{m!}$  centralizes  $\mathcal{E}(g^{m!})$ , hence  $\mathcal{E}(g^{m!}) = \{1\}$  in fact, so  $g^{m!}$  is an Engel element.

By Baer's theorem, then  $g^{m!} \in F(G)$ , so  $G/F(G)$  has exponent dividing  $m!$ . □

## Further proof for finite groups

### Proposition

*If  $\forall |\mathcal{E}(g)| \leq m$ , then  $|G/F(G)|$  is  $m$ -bounded.*

First for the case of soluble  $G$ .

Then considering the generalized Fitting subgroup = socle of  $G/S(G)$   
(using CFSG).....

**Proof of Theorem 2** (that  $|\gamma_\infty(G)|$  is  $m$ -bounded)

is by induction on  $|G/F(G)|$ ...

# Profinite groups

Recall:

Inverse limits of finite groups.

Topological groups. Quotients only by closed subgroups.

Open subgroups have finite index and are also closed.

Sylow theory. Pronilpotent (=pro-(finite nilpotent)) groups are Cartesian products of pro- $p$  groups.

Largest normal pronilpotent subgroup (closed).

## Lemma

*A pronilpotent almost Engel group  $H$  is in fact an Engel group.*

**Proof:** For any  $h \in H$  there is a normal subgroup  $R$  such that  $\mathcal{E}(h) \cap R = \{1\}$  with nilpotent  $H/R$ .

Then  $\mathcal{E}(h) \subseteq R$ , so in fact  $\mathcal{E}(h) = \{1\}$ ,

so  $h$  is an Engel element. □

## Bounded version for profinite groups

Theorem 2 on finite groups immediately implies the following.

### Corollary

*Suppose that  $G$  is an almost Engel profinite group and there is a positive integer  $m$  such that  $|\mathcal{E}(g)| \leq m$  for every  $g \in G$ .*

*Then  $G$  has a finite normal subgroup  $N$  of order bounded in terms of  $m$  such that  $G/N$  is locally nilpotent.*

## General case of profinite groups

### Theorem 3

*Suppose that  $G$  is an almost Engel profinite group. Then  $G$  has a finite normal subgroup  $N$  such that  $G/N$  is locally nilpotent.*

Cannot simply apply Theorem 2 on finite groups – as there is no a priori uniform bound on  $|\mathcal{C}(g)|$ .

First goal: a pronilpotent normal subgroup of finite index.

In the proof, a certain section is considered, and the Baire category theorem is applied.

# A piece of proof

## Lemma

In an almost Engel profinite group  $G$ , the sets

$$E_k = \{x \mid |\mathcal{E}(x)| \leq k\}$$

are closed.

**Proof:** For  $y \notin E_k$  we have  $|\mathcal{E}(y)| \geq k + 1$ , so there are  $z_1, z_2, \dots, z_{k+1}$  distinct elements, each

$$z_i = [z_i, y, \dots, y]. \quad (*)$$

There is an open normal subgroup  $N$  such that the images of the  $z_i$  are distinct in the finite quotient  $G/N$ .

Then equations (\*) show that for every  $n \in N$  the sink  $\mathcal{E}(yn)$  has an element in every coset  $z_i N$ , whence  $|\mathcal{E}(yn)| \geq k + 1$ . So  $yN$  is also contained in  $G \setminus E_k$ . Thus,  $G \setminus E_k$  is open, so  $E_k$  is closed. □

## Application of Baire theorem

Recall:  $E_k = \{x \mid |\mathcal{E}(x)| \leq k\}$  are closed.

In the theorem,  $G$  is almost Engel, which means  $G = \bigcup E_k$ .

By the Baire category theorem, one of  $E_k$  contains an open set, coset  $aU$ , where  $U$  is an open subgroup.

This gives us, in a certain metabelian section, a uniform bound for  $|\mathcal{E}(u)|$  for all  $u \in U$ , and then Theorem 2 on finite groups can be applied...

Thus,  $|G/F(G)|$  is finite, where  $F(G)$  is the largest pronilpotent normal subgroup (which is also locally nilpotent by Lemma above). Further arguments are by induction on  $|G/F(G)|$  and are similar to those for finite groups.

# Compact groups

Recall

## Theorem 1

*Suppose that  $G$  is an almost Engel compact group. Then  $G$  has a finite normal subgroup  $N$  such that  $G/N$  is locally nilpotent.*

## Structure theorems for compact groups:

- The connected component  $G_0$  of the identity is a divisible group (that is, for every  $g \in G_0$  and every integer  $k$  there is  $h \in G_0$  such that  $h^k = g$ ).
- $G_0/Z(G_0)$  is a Cartesian product of simple compact Lie groups.
- $G/G_0$  is a profinite group.

Note that [a simple compact Lie group is a linear group](#).

## $G_0$ is abelian

### Lemma

*An almost Engel divisible group is an Engel group.*

**Proof:** For  $g \in G_0$ , let  $|\mathcal{E}(g)| = m$ . Choose  $h \in G_0$  such that  $h^m = g$ . Clearly,  $h$  centralizes  $g$ , so for any  $z \in \mathcal{E}(g)$  we have

$$z = [z, g, \dots, g] \Rightarrow z^h = [z^h, g, \dots, g].$$

Hence  $\mathcal{E}(g)$  is  $h$ -invariant. Then  $h^m = g$  centralizes  $\mathcal{E}(g)$ . This means that actually  $\mathcal{E}(g) = \{1\}$ , so  $g$  is an Engel element. □

By the structure theorem,  $G_0$  is divisible, so is Engel by the above.

By well-known results (Garashchuk–Suprunenko, 1960), linear Engel groups are locally nilpotent.

Hence  $Z(G_0) = G_0$  is abelian by the structure theorem.

## Using the profinite case

We apply Theorem 3 on profinite groups to  $G/G_0$ .

Thus we have  $G_0 < F < G$  with  $G_0$  abelian divisible,  $F/G_0$  finite, and  $G/F$  locally nilpotent.

Next steps:

$\mathcal{E}(g) \cap G_0 = \{1\}$  for all  $g \in G$ ;

$[G_0, \mathcal{E}(g)] = 1$  for all  $g \in G$ ;

Replace (rename)  $F$  by possibly smaller subgroup  $\langle \mathcal{E}(g) \mid g \in G \rangle G_0$ ,  
so  $G_0 \leq Z(F)$ ;

... etc., in the end use Theorem 3 on profinite again.

## Almost Engel in the sense of rank

Instead of being finite, suppose that  $\mathcal{E}(g)$  generates a subgroup of finite (Prüfer) **rank**, for all  $g \in G$ .

### Conjecture:

If  $G$  is a compact (or profinite) group, then there is a normal closed subgroup  $N$  of finite rank such that  $G/N$  is locally nilpotent.

So far, the case of finite groups has been done:

### Theorem 4

*Suppose that  $G$  is a finite group and there is a positive integer  $r$  such that  $\langle \mathcal{E}(g) \rangle$  has rank at most  $r$  for every  $g \in G$ . Then the rank of  $\gamma_\infty(G)$  is bounded in terms of  $r$ .*